

CARROLL COUNTY: Key Data Points

- Carroll county's total birth rate is slightly lower than the state of Indiana overall, 11.6 versus 13.9 respectively.
- The percentage of women who smoked during pregnancy is lower than that of the state of Indiana (16.5% versus 18.5% respectively), as well as the percentage of unmarried parents, 29.5% versus 37.1% respectively.
- Carroll county observed 14 accidental deaths (4 of which were motor vehicle) and 1 suicidal death.

Note: All figures relate to 2003 data from the Indiana State Department of Health, unless otherwise specified

Figure 1. Births

	Total		White		Black		Other	
	Number	Crude Rate*	Number	Crude Rate*	Number	Crude Rate*	Number	Crude Rate*
Carroll County	237	11.6	234	11.5	0	0(U)	3	34.9(U)
INDIANA	86,382	13.9	75,422	13.7	9,288	17.5	1,672	10.6

* births per 1,000 population; U = unstable rate, based on fewer than 20 births

Figure 2. Mortality

		NUMBER				RATE*		
COUNTY OF RESIDENCE	REPORTED RACE/ ETHNICITY	Live Births	Infant Deaths (< 1 yr)	Neonatal Deaths (< 28 days)	Post Neonatal Deaths (28-364 days)	Infant Deaths (< 1 yr)	Neonatal Deaths (< 28 days)	Post Neonatal Deaths (28-364 days)
CARROLL	All Races	237	0	0	0	0	0	0
	White	234	0	0	0	0	0	0
	Black	0	0	0	0	0	0	0
	Other	3	0	0	0	0	0	0
	Hispanic	16	0	0	0	0	0	0
INDIANA	All Races	86,382	641	417	224	7.4	4.8	2.6
	White	75,422	479	309	170	6.4	4.1	2.3
	Black	9,288	148	98	50	15.9	10.6	5.4
	Other	1,672	14	10	4	**	**	**
	Hispanic	6,763	38	22	16	5.6	3.3	**

*rate per 1,000 live births **Numerator is less than 20 and the rate is unstable

Figure 3. Outcome Indicators of Mother, by Race

		Birth Weight				During Pregnancy		
CARROLL COUNTY	Total Births	% Low (< 2500 grams)	% Very Low (< 1500 grams)	% Preterm †	% PNC 1st Trimester ‡	% Alcohol	% Smoked	% Unmarried Parents
All Races	237	8.4	1.3	7.6	85.7	0.4	16.5	29.5
White	234	8.5	1.3	7.7	86.3	0.4	16.7	29.1
Black	0	**	**	**	**	**	**	**

		Birth Weight				During Pregnancy		
INDIANA	Total Births	% Low (< 2500 grams)	% Very Low (< 1500 grams)	% Preterm †	% PNC 1st Trimester ‡	% Alcohol	% Smoked	% Unmarried Parents
All Races	86,382	7.9	1.4	9.9	80.6	0.6	18.5	37.1
White	75,422	7.2	1.2	9.5	82.1	0.6	19.1	32.6
Black	9,288	13.3	3.3	13.3	68.2	1.1	15.2	76.3

† Preterm: less than 37 weeks gestation

‡ PNC: Prenatal Care

'S' Significantly different than the state percent

'**' Outcomes for this county are suppressed due to 20 or fewer total births

Figure 4. Breastfeeding

CARROLL COUNTY	All Races	White	Black
Total Births	237	234	0
Breastfeeding on Discharge	160	157	0
% Breastfeeding on Discharge	67.5	67.1	--




Figure 4. Breastfeeding

INDIANA	<i>All Races</i>	<i>White</i>	<i>Black</i>
Total Births	86,382	75,422	9,288
Breastfeeding on Discharge	54,636	49,385	3,959
% Breastfeeding on Discharge	63.2	65.5	42.6




Figure 5. HIV/AIDS

	Total # of AIDS cases reported since 1982	Total # of HIV cases reported since 1985
CARROLL COUNTY	4	1
INDIANA	7,641	3,856

*data gathered from June 2005 HIV/STD Quarterly Report

Figure 6. Nutritional Risk Factors by Status,
Indiana WIC Clinics, Quarter Ending 09/30/2004

Local Agency: White County Memorial Hospital							
Counties Included: White, Carroll							
Risk Factor or Status	Children 1-5 years of Age					Total	% within Agency
	White	Black	Hispanic	Asian	Other		
Overweight	8	0	7	0	0	15	3.6%
At Risk for becoming Overweight	12	0	9	0	1	22	5.2%
Local Agency: State Wide							
Counties Included: All							
Risk Factor or Status	Children 1-5 years of Age					Total	% within Agency
	White	Black	Hispanic	Asian	Other		
Overweight	1,732	445	800	12	97	3,086	4.8%
At Risk for becoming Overweight	2,023	485	639	11	124	3,282	5.1%

Priority Issues (Carroll & Indiana)			
Demographics: Percentage of population enrolled in Medicaid, Percentage of population enrolled in Food Stamps.		Carroll	Indiana
2	% of Population enrolled in Medicaid as of June 2003	7.7	12.4
3	% of Population in Food Stamps Program, 2004	4.3	8.6
Priority #1 (pregnancy outcomes): Average Infant Mortality Rate, Average Fetal death Rate		Carroll	Indiana
1	Av Total Infant Death Rate (1993-2002)	NS	8.1
2	Average Fetal Death Rate (1994-2003)	4.5	6.7
Priority #2 (reduce barriers): Health Professional Shortage Areas, Percentage of women with prenatal care in first trimester		Carroll	Indiana
1	Scores for Primary Care Health Professional Shortage Areas (HPSAs) (2005)	No Score	NA
2	% PNC 1st Trimester†, 2003	85.7	80.6
Priority #4 (environmental): Prevalence of asthma, Childhood lead poisoning		Carroll	Indiana
1	Prevalence of Asthma in a Medicaid Cohort of Children age 0-17 years, 2003	10.5	10.4
2	%Estimated Lead Blood Levels (%EBLL) for Children under 6 years old FY 2004-2005	NA	2.1
Priority #5 (tobacco): Percentage of mothers who smoked during pregnancy, Percentage of Adult Smoking		Carroll	Indiana
1	% of mothers Smoked During Pregnancy, 2003	16.5	18.5
2	% of Adult Smoking in 2003	29.9	29.3
Priority #7 (adolescent risk behaviors): Adolescent deaths due to motor vehicle accidents, Chlamydia rates (ages 15-24).		Carroll	Indiana
1	Number of Adolescent (15-24) deaths due to Motor vehicle accidents (2001-2003)	4	748
2	Chlamydia rates per 100,000 county specific population (15-24yrs of Age)	296.85	1541.49
Priority #8 (obesity): Obesity in pregnant women		Carroll	Indiana
1	Number of Obese Pregnant Women In Indiana, 2003	3	2,496
Priority #9 (domestic violence, childhood injury): Child deaths due to Injury, Child abuse and neglect rates		Carroll	Indiana
1	Child deaths due to Injury 1-14yrs (2001-2003)	2	257
2	Child Abuse and Neglect per 1,000 Population under age 18, 2004	10.8	13.5
Priority #10 (To improve racial and ethnic disparities in women of childbearing age, mothers, and children's health outcomes): Percentage of Hispanic Population of any Race, LBW Disparity Ratio		Carroll	Indiana
1	% of Hispanic Population of any Race, July,1,2004	3.6	4.3
2	Low Birth Weight Disparity Ratio (B/W) (1993-2002)	NS	1.92
Source: Data Analysis Team, ISDH. Notes: 1. NS= Not Stable 2. NA = Not Available 3. Infant Mortality rate and Fetal Death rate is for 1000 live births. 4. Low Birth Weight: Birth Weight Less than 2,500grams. 5. HPSA Score ranges 1 – 25. The higher the score, the higher the need. 6. EBLL: Ten (10) micrograms of lead per deciliter of whole blood for one venous test; %EBLL is based on screening rate of 27% Statewide, unduplicated kids 7. Obesity in Pregnant Women determined by "Concurrent Illness Obesity" checkbox on the birth certificate 8. PNC percentages are calculated using total births in each county.			

